



Research Article

At-Sea Distribution of Seabirds and Marine Mammals in the Unexplored Southern Weddell Sea, Antarctica

Claude R. Joiris , Diederik D’Hert

Laboratory for Polar Ecology (PoE), Belgium

KEYWORDS

seabirds & marine mammals at-sea distribution
geographical heterogeneities
Weddell Sea
S. georgia, S. Sandwich
S Shetlands islands
antarctic continent

ABSTRACT

Background: In the frame of our long-term study in polar seas, the at-sea distribution of “top predators” – seabirds and marine mammals – was studied in the Weddell Sea area, both the well-covered northern one (sub-antarctic Scotia Sea) as well as in the almost unexplored southern and central ones. Method was our usual transect counts lasting 30 min. each, without width limitation. **Results and conclusions:** Species richness was low, dominated by only a few species, reflecting low diversity. Birds, seals and to some extent whales showed a heterogeneous distribution. They were concentrated in a few “hotspots” on the “continental” shelf and –slope, of South Georgia, South Sandwich and South Shetland (Elephant) islands, and off the Peninsula. Species showed important differences in distribution probably reflecting differences in the food web. In open waters off the shelf, species showed low densities, both in the northern and southern areas. In the case of seabirds, moreover, they were partially over-estimated by the presence of long-distance followers, probably by one order of magnitude. The absence of gentoo penguin might reflect a change in geographical distribution.

*CORRESPONDING AUTHOR

Claude R. Joiris; Laboratory for Polar Ecology (PoE), Belgium; Email: crjoiris@gmail.com

ARTICLE INFO

Received: 27 January 2026 | Revised: 13 February 2026 | Accepted: 18 February 2026 | Published Online: 25 February 2026

DOI: <https://doi.org/10.65773/pcrr.2.1.114>

COPYRIGHT

Copyright © 2026 by the author(s). Published by Explorer Press Ltd. This is an open access article under the Creative Commons Attribution 4.0 International (CC BY 4.0) License (<https://creativecommons.org/licenses/by/4.0>)

1. Introduction

Our study of the at-sea distribution of the marine “top predators” in polar seas consists in a synecological approach providing information on the functioning of the whole ecosystem. The number of species can reflect biodiversity, while the number of individuals can indicate productivity (prey abundance). Species richness is generally low in polar seas – approximately 30 bird species – with high heterogeneity in distribution. The activity of the Laboratory for Polar Ecology (PoE) team started with a pioneering return expedition from Belgium to Greenland in 1973 [1] and was followed by many others in both polar areas. In the context of our long-term study of the at-sea distribution of seabirds and marine mammals in polar seas, we had the opportunity to cover an almost unexplored area of the southern Weddell Sea.

2. Method

An observer of the PoE team (CJ) participated in a cruise on a small tourist ship (90 m long, 80 passengers), into the Scotia and Weddell seas, from Ushuaia, Argentina, to the Antarctic continent, from 12 February to 9 March 2025. The vessel was not an icebreaker, but ice strengthened, and so it avoided areas of heavy sea ice cover. Seabirds and marine mammals were counted by line transect methodology, on a continuous basis, visibility conditions allowing. Continuous transects were broken into 30 min segments without width limitation. Data were collected from one side and in front of the bridge in function of light conditions, 9 m above the sea-level and were facilitated by using 10X40B binoculars when necessary. As a consequence of the low numbers of individuals in polar areas, the counting methodology had to be adapted, not following the “compulsory” method of 10 min transect counts with a width limitation of 300 m adapted for areas with very high local abundance such as the northern Atlantic Ocean - besides there exists no compulsory methodology in any science. In order to compare with data from previous expeditions by the same team and same methodology, mean values were used to correct for the number of counts instead of medians – even if the distribution is far from normal with median values of zero, but for the sum of all birds equal to seven.

3. Results and Discussion

3.1. Tallies of Birds and Mammals

The cruise and its 357 30 min transect counts basically covered the northern and the almost unexplored southern areas of the Weddell Sea (Figure 1), as compared with the radar-registered ships for decennia (Figure 2). The southwestern portion of the Weddell Sea had to be avoided due to heavy ice coverage (Figure 3). In total, 30 seabirds, five pinnipeds and eight cetaceans were encountered, dominated by a few species, typical of polar seas (Table 1). Some hotspots on the continental plateau and –slope presumably reflected high food availability, the reasons for which are not known. High concentrations were observed mainly in four areas, off S. Georgia, S. Sandwich and S. Shetland (Elephant) islands and the Peninsula and to a lower extend off the antarctic continent (Figure 4). Species distributions tended to be heterogeneous (Figure 5). For instance, 564 fulmars were counted sitting on ice floes or on water next to ice floes in seven counts in the same area off the Peninsula and Elephant Island (Photo 1), out of the total of 940 for the whole cruise. The most abundant species, antarctic prion and blue petrel, were observed in very large flocks flying back to their breeding colonies and thus do not reflect food web generated concentrations.



Photo 1. Flocks of fulmars sitting on, or close to, medium ice floes




Figure 1. Position of the seabirds and marine mammals 30 min line transect counts in the Weddell Sea, February-March 2025.




Figure 2. Route of this cruise (thin line) compared with the radar-registered ships in the area (colours).




Figure 3. Heavy ice coverage in the SW Weddell Sea (red).




Figure 4. Sum of the observations of all seabirds (a), all pinnipeds (b) and all cetaceans (c); numbers per count.

Hotspots were embedded in empty zones (many counts had no detection). High counts in open waters otherwise having low bird numbers often involved species that tend to follow vessels, causing an overestimation of density, probably by one order of magnitude: e.g. giant petrel, black-browed albatross, pintado (Cape), snow and antarctic petrels. Two areas stood out: a northern one that included the islands and Peninsula, north of S Sandwich Islands and the Peninsula, and a southern one that was partially covered with medium ice floes and many icebergs..

Among the most abundant penguin species were emperor and Adélie, which were mostly seen on old icebergs and ice floes. Other bird species were clearly bound to the island coasts. We note that numbers of emperors were artificially higher, owing to attraction by tens around the stationary ship, being curious and attracted by people walking on the ice or by Zodiac boats and divers close to the ice floes (Photo 2). All hotspots were situated around coasts above the continental shelf and –slope (depths of 300 to 400 m).

Table 1. Seabirds and marine mammals encountered at sea in the Weddell Sea, February-March 2025.

	Zone*	All		Norh		South		Remark
		Total	Mean**	Total	Mean*	Total	Mean**	
Species***	Number of counts	357		195		157		
1 king penguin	<i>Aptenodytes patagonicus</i>	47	0.13	47	0.24	0		
2 emperor penguin	<i>Pygoscelis papua</i>	129	0.36	93	0.48	36	0.24	
gentoo penguin	<i>Pygoscelis papua</i>	0		0		0		
3 chintrap penguin	<i>Pygoscelis antarctica</i>	15	0.06	15	0.08	0		and 2 large colonies on Elephant Isl
4 Adélie penguin	<i>Pygoscelis adeliae</i>	115	0.32	33	0.17	82	0.52	
5 rockhopper penguin	<i>Eudyptes chrysocome</i>	6		6		6		off Ushuaia
penguin sp		51	0.14	45	0.23	6		
6 wandering albatross	<i>Diomedea [exulans] sp</i>	42	0.14	42	0.22	0		
7 royal albatross	<i>Diomedea [epomorpha] epomorpha</i>	30	0.12	27	0.14	3		
8 black-browed alb.	<i>Thalassarche [melanophrys] melanophrys</i>	105	0.29	104	0.53	1		
9 grey-headed albatross	<i>Thalassarche chrysostoma</i>	7		7		0		
10 sooty albatross	<i>Phoebastria fusca</i>	10	0.03	0		10		
11 light-mantled albatross	<i>Phoebastria palpebrata</i>	3		3		0		
12 giant petrel sp	<i>Macronectes sp</i>	407	1.15	358	1.84	49	0.31	both N and S species
13 antarctic petrel	<i>Thalasoica antarctica</i>	283	0.80	152	0.78	133	0.84	peninsula
14 pintado (Cape) petrel	<i>Daption capense</i>	210	0.59	195	1.00	20	0.13	
15 snow petrel	<i>Pagodroma [nivea]</i>	435	1.22	268	1.37	149	0.94	
16 blue petrel	<i>Halobaena caerulea</i>	1528	4.30	83	0.43	1445	9.15	
17 southern fulmar	<i>Fulmarus glacialis</i>	941	2.65	909	4.66	33	0.21	
18 (antarctic) prion	<i>Pachyptila (desolata)</i>	1674	4.73	1360	7.01	314	1.99	
19 Kerguelen petrel	<i>Pterodroma brevirostris</i>	12	0.03	12	0.06	0		
20 soft-plumaged petrel	<i>Pterodroma mollis</i>	174	0.49	161	0.83	13	0.08	
21 white-chinned petrel	<i>Procellaria aequinoctialis</i>	198	0.56	198	1.03	0		off Ushuaia
22 sooty shearwater	<i>Puffinus griseus</i>	19	0.05	4	0.02	15	0.09	
23 Wilson storm-petrel	<i>Oceanites oceanicus</i>	319	0.89	300	1.54	19	0.12	
black-bellied storm-petrel	<i>Fregetta tropica</i>	46	0.13	41	0.21	5		mainly Peninsula
24 (georgean) diving-petrel	<i>Pelecanoides georgicus</i>	7		6		1		
26 south georgian shag	<i>Phalacrocorax (atriceps) georgianus</i>	26	0.07	25	0.13	0		
27 rock shag	<i>Phalacrocorax magellanicus</i>	125	0.35	125	0.64	0		off Ushuaia
28 (antarctic) tern	<i>Sterna vittata</i>	22	0.03	17	0.08	5		
29 (antarctic) skua	<i>Catharacta [skua] antarctica</i>	12	0.03	11	0.06	1		
total seabirds		6998	19.6	4647	23.8	2340	14.9	
snowy sheatbill	<i>Chionis alba</i>	10		9		1		terrestrial
1 (antarctic) fur seal	<i>Arctocephalus (pusillus) doriferus</i>	140	0.39	139	0.71	1		
2 leopard seal	<i>Hydrurga leptonyx</i>	7		5		2		

3	Weddell seal	<i>Leptonychotes weddellii</i>	4		2		2		
4	crabeater seal	<i>Lobodon carcinophaga</i>	121	0.34	120	0.61	1		
5	S elephant seal	<i>Mirunga leonina</i>	9		9		0		
	seal sp		21	0.07	20	0.10	1		
	total pinnipeds		299	0.85	292	1.52	7		
1	dolphin sp	<i>Lagenorhynchus</i> sp	32	0.99	32	0.16	0		probably Hourglass, 1 possible Peale's
2	long-finned pilot whale	<i>Globicephala melas</i>	13	0.03	9		4		
3	southern right whale	<i>Eubalena australis</i>	2		2		0		
4	antarctic Minke whale	<i>Balaenoptera bonaerensis</i>	31	0.08	6		25	0.15	
5	sei whale	<i>Balaenoptera borealis</i>	25	0.07	25	0.13	0		1 pod off Ushuaia
6	blue whale	<i>Balaenoptera (musculus) intermedia</i>	2		1		1		
7	humpback whale	<i>Megaptera novaeangliae</i>	47	0.13	20	0.10	27	0.17	
8	fin whale	<i>Balaenoptera physalis</i>	12	0.03	10	0.05	2		
	large whale sp		127	0.37	73	0.38	54	0.35	long-distance blows
	total cetaceans		312	0.88	200	1.04	112	0.71	

* zones: see table 1


** mean per count, when more than 10 individuals in total

*** species name in brackets: not all individuals were identified at the species level.



Photo 2. Emperor penguins attracted by the stationary ship and by people on and around pack ice, out of effort thus.

Cetaceans exhibited a distribution similar to that of seabirds, with humpback encountered somewhat widely and fin whale in the north. Among pinnipeds, the most numerous were antarctic fur seal and crabeater and Weddell seals along the coast, resting on ice floes (Table 1; Figure 5).






Figure 5. Detailed distribution of selected abundant species: emperor penguin (a), Adélie penguin (b), southern fulmar (c); giant petrel (d); blue petrel & prion (e); fur seal (f); numbers per count.

3.2. Comparison with Previous Expeditions in the Same Area

In comparison with previous expeditions of the same team in the same area, we used the data obtained in the Scotia Sea portion of this cruise and the Weddell Sea. The highest concentrations were, by far, observed around the S. Shetland Islands and Peninsula, with thousands of Adélie and chinstrap penguins, but also black-browed and grey-headed albatrosses, antarctic prions, or fur seal (Table 2).

Lower values for dolphins and penguins: possibly a seasonal effect, with the end of the breeding season and/or a different winter distribution, even if the seasons were similar: March-April 2012 and 2013. Reproducibility of data collected on successive, identical routes is known to be low – by up to one order of magnitude [2-4]. This is related to the heterogeneity of the distributions: the “hotspots” detected in both antarctic and arctic seas reflect the heterogeneities in prey abundance [5] (synthesis in [6]). No gentoo penguins were encountered, while registered at high concentrations during the previous expeditions in the same area: 0.2 and 0.4 per count (see Table 2).

A broader discussion consists in comparing data collected elsewhere in the Antarctic seas by the same team and same methodology. Concerning the number of species, data are of the same order in the different areas, reflecting similar species diversities. Densities (individuals per count) vary: less seabirds in this area, similar densities for pinnipeds, higher for cetaceans (Table 3). Such differences are considered to reflect differences in food availability and thus bio-productivity (see also [2,7-10]).

Table 2. Seabirds and marine mammals counted along the transects to S Georgia (A), S Shetlands (B) and S Sandwich islands (C) by tis team, mean number per 30 min transect count.

Transect	A	B	C
Number of counts	333	365	143
Species			
gentoo penguin	0.50	0.65	-
Adélie penguin	10.6	-	0.34
chinstrap penguin	14.3	7.81	0.06
black-browed albatross	4.23	2.01	0.45
grey-headed albatross	20.1	0.14	0.04
antarctic fulmar	231	1.94	0.73
antarctic prion	30.7	27.0	11.1
white-chinned petrel	0.31	2.56	0.57
sooty shearwater	1.33	0.18	-
giant petrel	1.43	1.66	2.04
Wilson storm-petrel	0.48	0.93	1.98
pintado (Cape) petrel	5.54	0.09	1.22
black-bellied storm-petrel	4.82	2.16	0.31
∑ birds	341	57	22
antarctic fur-seal	0.34	4.45	0.77
∑ pinnipeds	0.34	6.25	0.91
Dolphin sp	0.71	0.26	0.03
humpback whale	0.01	0.11	0.13
fin whale	0.15	0.02	0.05
∑ cetaceans	0.19	0.50	1.02

- A. Return transects to S Georgie and S Shetland isl [3]
- B. Return transects to S Georgia and S Sandwich isl [1]
- C. To S Georgia and S Sandwich isl, this study, partim

3.3. Comparison with Other Antarctic Seas

Table 3. Summary of the seabird and marine mammal at-sea distribution in Antarctic seas: main species.

Area	Seabirds			Pinnipeds		Cetaceans		References
	Number of counts	Number of species	Mean nbr per count	Number of species	Mean nbr per count	Number of species	Mean nbrper count	
Weddell Sea pack ice	291	24	180	6	6	3	0,1	[20]
Africa - Antarctica transects	1927	27	35	7	0,4	3	0,7	[3,4]
Amundsen & Bellingshausen seas	1000	15	8	4	2,3	5	0,2	[21]
off antarctic peninsula	100	25	40	3	7	3	0,5	Joiris, unpublished
S Georgia / S Sandwich transect	365	34	57	3	7	8	0,5	[21]
S Weddell Sea (northern area)	195	29	20	5	0,9	8	0,9	Joiris & D'Hert, this study
Mean		25.7	56,7	4,7	3,9	5	0,5	

4. Conclusion

The northern area, constituting Ushuaia to S. Georgia, S. Sandwich and S. Shetland Islands/ Peninsula (subantarctic Scotia Sea and Weddell Sea), corresponds to an area with local high densities of top predators [2,11-16] and prey availability [5,17]. The concentration of birds and mammals in the Scotia-Weddell Confluence, was already described [18,19].

Numbers of species was low, with highest numbers of individuals represented by few species. The distribution of seabirds, seals and to some extent cetaceans were heterogeneous and were mainly observed above shallow areas along the coasts, corresponding to the continental slope of the islands and the continent: this is consistent with these areas generally characterized by high prey availability. The southern Weddell Sea, almost unexplored, showed in contrast very low densities in open water, among the lowest of the Antarctic seas, even if partially over-estimated by the presence of long-distance followers in low numbers, registered once per count.

Contributions

Claude R. Joiris on board and basic redaction; Diederik D'Hert maps. Data set available by Claude R. Joiris

Funding

No external funding

Conflicts of Interest

The authors don't have any conflicts of interest.

Acknowledgments

We are grateful to D. Ainley, J.-M. Valette and two anonymous reviewers who helped improve a first version of this manuscript.

References

- [1] Joiris C. 1976. Seabirds seen during a return voyage from Belgium to Greenland in July. *Gerfaut* 66, 63-87.
- [2] Joiris CR. & Dochy O. 2013. A major autumn feeding ground for fin whales, southern fulmars and grey-headed albatrosses around the South Shetland Islands, Antarctica. *Polar Biol.* 36, 1649-1658. <https://doi.org/10.1007/s00300-013-1383-8>.
- [3] Joiris CR, Humphries GRW, D'Hert D et al. 2015. Major hotspots detected along the Scotia Ridge in autumn for southern right whales *Eubalaena australis*, Antarctic fur seals *Arctocephalus gazella* and Antarctic prions *Pachyptila desolata*. *Adv. Polar Sc.* 26, 282-291. <https://doi.org/10.13679/j.advps.2015.4.00282> <http://www.aps-polar.org/paper/2015/26/04/A20150403>.
- [4] Joiris CR. 2021a. On the reproducibility of bird and mammal transect counts in polar seas. *J Mar Sci Res Ocean* 4, 145-153. <https://doi.org/10.33140/JMSRO.04.01.01>.
- [5] Veit RR, Silverman ED & Everson I. 1993. Aggregation patterns of pelagic predators and their principal prey, antarctic krill, near South Georgia. *J Animal Ecol.* 62, 551-564. <http://www.jstor.org/stable/5204>.
- [6] Joiris C. 2018. Seabird and marine mammal "hotspots" in polar seas. Lambert Academic Publishing, Düsseldorf, Germany. 48 ps. ISBN: 978-620-2-19805-9. For sale: morebooks.de., amazon, 35.9€.

- [7] Ainley DG, Jacobs S, Ribic CA, & Gaffney I. 1998. Seabird distribution and oceanic features of the Amundsen and southern Bellingshausen seas. *Antar Sci* 10, 111-123.
- [8] Ainley DG, Dugger KM, Toniolo V, & Gaffney I. 2007. Cetacean occurrence patterns in the Amundsen and southern Bellingshausen Sea sector, Antarctic seas. *Mar Mamm Sci* 23, 285-305.
- [9] Bassoi M, Acevedo J, Secchi ER, et al. 2020. Cetacean distribution in relation to environmental parameters between Drake Passage and northern Antarctic Peninsula. *Polar Bol.* 43 preprint; <https://doi.org/10.1007:s00300-019-02607-z>.
- [10] Joiris CR. 2020. Abundance and biodiversity of top predators – seabirds and marine mammals – in Antarctic seas. *J Mar Sci Res Ocean* 3, 89-99. <https://doi.org/10.33140/JMSRO.03.03.07>
- [11] Jackson JA, Kennedy A, Moore M et al.. 2020. Have whales returned to a historical hotspot of industrial whaling? The pattern of southern right whale *Eubalaena australis* recovery at South Georgia. *Endang Species Res.* 43, 323-339. <https://doi.org/10.3354/esf01072>.
- [12] Chernetsky AD, Chukmasov PV, Krasnova VV et al. 2023. Occurrence and Distribution of Fin Whales (*Balaenoptera physalus*) and Humpback Whales (*Megaptera novaengliaei*) in the Atlantic Sector of Antarctica. *Oceanology* 63, 589-599 [from a Russian text in *Okeanologiya* 63, 677-688].
- [13] Calderan SV, Dornan TJ, Fielding S. et al. 2023. Observations of southern right whales (*Eubalaena australis*) surface feeding on krill in austral winter at South Georgia. *Marine Mam Sci* 39. <https://doi.org/10.1111/mms.13025>.
- [14] Owen KA, Goggin S. L, Black A et al. 2024. At-sea distribution of marine predators around South Georgia during austral winter, with implications for fisheries management. *Polar Biol.* 47, 663-679. <https://doi.org/10.1007/s00300-024-03257-6>.
- [15] Rand ZR, Branch TA & Jackson JA. 2024. High historical movement rates of Antarctic blue whales on Antarctic seas feeding grounds estimated from Discovery mark data. *Endang. Species. Res.* 55, 109-128. <https://doi.org/10.3354/esr01361>.
- [16] Bitiutskii, DG, Samyshev EZ, Minkina NI et al. 2022. Distribution and Demography of Antarctic Krill and Salps in the Atlantic Sector of the Antarctic seas during Austral Summer 2021–2022. *Water* 14, 3812. <https://doi.org/10.3390/w14233812>.
- [17] Machado-Gaye AL, Mania G, Braun C et al., 2025. Divergent responses of *Pygoscelis* penguins to unfavourable weather conditions in the South Shetland Islands. *Antar. Sci* 37, 384-394. <https://doi.org/10.1017/S0954102025100291>
- [18] Hopkins TL, Ainley DG, Torres JJ & Lancaft TM. 1993. Trophic structure in open waters of the marginal ice zone in the Scotia-Weddell confluence region during spring (1983). *Polar Biol.* 13, 389-397.
- [19] Ainley D.G., Ribic CA & Fraser WR. 1994. Ecological structure among migrant and resident seabirds of the Scotia-Weddell Confluence region. *J. Anim. Ecol.* 63, 347-364.
- [20] Joiris CR. 1991. Spring distribution and ecological role of seabirds and marine mammals in the Weddell Sea, Antarctica. *Polar Biol.* 11, 415-424.
- [21] Joiris CR. 2021b. Very low density and biodiversity of seabird and marine mammal distribution in the Amundsen Sea, Antarctica, 2010. *J Mar Scie Res Ocean*, 4, 198-204. <https://doi.org/10.33140/JMSRO.04.02.02>.